



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

CALCULUS.

254. Proposed by H. S. PARDEE, Boston, Mass.

A wire is wound in the form of a helix. Assuming that sections of the wire perpendicular to the axis of the wire are circles, find the equation of a section of the wire perpendicular to the axis of the helix.

MECHANICS.

214. Proposed by W. J. GREENSTREET, M. A., Marling School, Stroud, England.

An inelastic particle is projected in a direction BD from B in a straight line AB . It strikes a rigid line AD in D and returns to AB at C . Find AC/AB , and show on *a priori* ground that this ratio is independent of the velocity of projection.

NUMBER THEORY AND DIOPHANTINE ANALYSIS.

152. Proposed by H. S. VANDIVER, Bala, Pa.

When p is a prime of the form $5n \pm 1$ then there is a positive integer a such that $a^2 \equiv 5 \pmod{p}$. Show that $\left(\frac{a \pm 1}{p}\right) = \pm \left(\frac{-2a}{p}\right)$, according as p is of the form $5n+1$ or $5n-1$.

AVERAGE AND PROBABILITY.

195. Proposed by G. B. M. ZERR, A. M., Ph. D., 4243 Girard Avenue Philadelphia, Pa.

A random diameter is drawn in a given circle. Find the chance that it intersects, (1) a random chord; (2) a random chord through a random point; and (3) a chord through two random points.

NOTES AND NEWS.

We learn from the *Scientific American* of February 1, 1908, that Dr. Paul Wolfskehl, who died recently at Darmstadt, left in his will a provision for the payment of 100,000 marks to the first person who will prove or disprove "the last theorem of Fermat," viz., $x^n + y^n = z^n$, is not possible in integers for $n > 2$. F.

The Fourth International Congress of Mathematicians will convene in Rome April 6th to 11th, 1908. Extensive preparations are announced for the entertainment of delegates and their friends. The deliberations of the Congress will be conducted under four sections, each provided with leaders of international reputation: (1) Arithmetic, Algebra, Analysis; (2) Geometry; (3) Mechanics, Mathematical Physics; (4) Philosophical, historical and didactic questions. Professor E. H. Moore, who is now sojourning in Italy, will represent The University of Chicago. S.